

# Retina Institute of Virginia

8700 Stony Point Parkway, Suite 150

Richmond, VA 23235

(804) 644-7478

Toll-Free (877) 348-3937

*Ali Tabassian, M.D., Ph.D., Medical Director*

*Juan A. Astruc, Jr., M.D.*

*J. Stewart O'Keefe, M.D.*

*Bradley T. Smith, M.D.*

## CARE AFTER YOUR SURGERY:

- Your operated eye will have a patch on after your surgery. Do not remove this yourself. It will be removed on your follow up appointment the following day.
- If you are given a wristband indicating you received a gas bubble during your surgery you must leave it on until you are instructed to take it off.
- After your 1<sup>st</sup> follow up appointment you may clean your eyelids gently with water and a clean cloth. You may shower or bathe, but DO NOT get soapy water in the operated eye.
- Continue to protect your eye as much as possible using your sunglasses provided in your post-op kit.
- Continue to use all medications as before your surgery. You may also take Tylenol or Ibuprofen for pain if you are not allergic to these medications.
- Don't drive the day of surgery or while taking narcotic pain medications.
- Do not drink alcoholic beverages for the next 24 hours or while taking narcotic medications.
- Do not make any important decisions or sign important papers for 24 hours.

\*\*\*If intense pain or nausea occurs, please contact our office as soon as possible.

## USE NEW EYE MEDICATIONS, AS FOLLOWS, AFTER YOUR FIRST FOLLOW UP VISIT UNTIL THEY ARE FINISHED:

- Econopred Plus 1% - One drop in the operated eye four times a day.
- Antibiotic drop – One drop in the operated eye four times a day.
- Use all other eye medications in the operated eye as before your surgery unless otherwise instructed.
- When using eye drops together you should space them apart by 5 minutes.
- Refill of drops is NOT necessary, unless otherwise instructed by your doctor and please bring all medications in with you to each appointment.

---

Patient/Representative Signature

---

MD/Nurse Signature

---

Date